# antibodies -online.com







# anti-CD19 antibody (AA 96-281)





### Overview

Quantity:	100 μg
Target:	CD19
Binding Specificity:	AA 96-281
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD19 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), ELISA, Coating (Coat), Staining Methods (StM)

#### **Product Details**

. Todast Dotalio		
Immunogen:	Recombinant fragment of human CD19 protein (around aa96-281) (exact sequence is proprietary)	
Clone:	CD19-3117	
Isotype:	IgG2b kappa	
Purification:	Purified by Protein A/G	
Target Details		

## l arget Details

Target:	CD19
Alternative Name:	CD19 (CD19 Products)

# **Target Details**

Storage:

rarger Details		
Background:	CD19 is a transmembrane glycoprotein that contains two extracellular immunoglobulin-like domains. CD19 is present in both benign and malignant B-cells and is considered to be the most reliable surface marker of this lineage over a wide range of maturational stages. In normal lymphoid tissue, CD19 is observed in germinal centers, in mantle zone cells, and in scattered cells of the inter-follicular areas. Anti-CD19 exhibits an overall immunoreactivity pattern similar to those of the antibodies against CD20 and CD22. However, in contrast to CD20, expression of CD19 is continuous throughout B-cell development and through terminal differentiation of B-cells into plasma cells. Anti-CD19 positivity is seen in the vast majority of B-cell neoplasms commonly at a lower intensity than normal B-cell counterparts. Plasma cell neoplasms are nearly always negative, as are T-cell neoplasms.	
Molecular Weight:	95kDa	
Gene ID:	930	
UniProt:	P15391	
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway	
Application Details		
Application Notes:	Positive Control: Raji cells (FACS). Tonsil, Lymph Node or Spleen (IHC).  Known Application: ELISA (For coating, order antibody without BSA), ,Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.	
Restrictions:	For Research Use only	
Handling		
Concentration:	200 μg/mL	
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	

4 °C,-80 °C


#### Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

#### **Images**




#### **SDS-PAGE**

**Image 1.** SDS-PAGE Analysis Purified CD19 Monospecific Mouse Monoclonal Antibody (CD19/3117). Confirmation of Purity and Integrity of Antibody.



#### **Protein Array**

Image 2. Analysis of Protein Array containing more than 19,000 full-length human proteins using CD19 Monospecific Mouse Monoclonal Antibody (CD19/3117). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SDs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SDs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



#### **Immunohistochemistry**

**Image 3.** Formalin-fixed, paraffin-embedded human Tonsil stained with CD19 Monospecific Mouse Monoclonal Antibody (CD19/3117).

Please check the product details page for more images. Overall 4 images are available for ABIN6941102.