

Datasheet for ABIN6941111

**Recombinant anti-CD20 antibody**

5 Images

[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	CD20 (MS4A1)
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This CD20 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF), Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant full-length human MS4A1 fragment
Clone:	RIGEL-773
Isotype:	IgG1 kappa
Specificity:	Recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed by pre-B-cells and persists during all stages of B-cell maturation but is lost upon terminal differentiation into plasma cells. This MAb can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue,

## Product Details

germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.

Purification: Purified by Protein A/G

## Target Details

Target: CD20 (MS4A1)

Alternative Name: MS4A1 ([MS4A1 Products](#))

Molecular Weight: 33-37kDa

Gene ID: 931

UniProt: [P11836](#)

## Application Details

Application Notes: Positive Control: Daudi, Raji or U266 (FACS). Human lymphocytes. Lymph node and tonsil (IHC).  
Known Application: Immunofluorescence (1-2 µg/mL), Flow Cytometry (1-2 µg/million cells), ,Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL


Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-80 °C

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.




### Protein Array

**Image 1.** Analysis of Protein Array containing more than 19,000 full-length human proteins using CD20 Monospecific Recombinant Mouse Monoclonal Antibody (rIGEL/773). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



### Flow Cytometry

**Image 2.** Flow Cytometric Analysis of Raji cells using CD20 Mouse Recombinant Monoclonal Antibody (rIGEL/773) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



#### Immunofluorescence

**Image 3.** Immunofluorescence staining of Raji cells using CD20 Mouse Recombinant Monoclonal Antibody (rIGEL/773) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN6941111.