

Datasheet for ABIN6941146  
**anti-TNFRSF8 antibody**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	TNFRSF8
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNFRSF8 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Human CD30 recombinant protein
Clone:	CD30-412
Isotype:	IgG1 kappa
Specificity:	Recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In Hodgkin's disease, CD30/Ki-1 antigen is expressed by mononuclear-Hodgkin and multinucleated Reed-Sternberg cells. It is also expressed by the tumor cells of a majority of anaplastic large cell lymphomas as well as by a varying proportion of activated T and B cells. This MAb distinguishes large cell lymphomas derived from activated lymphoid cells from histiocytic malignancies and lymphomas derived from resting and precursor lymphoid cells or from anaplastic carcinomas. About one third of the Ki-1 positive lymphomas lack the leukocyte common antigen (CD45).

## Product Details

---

Purification: Purified by Protein A/G

## Target Details

---

Target: TNFRSF8

Alternative Name: TNFRSF8 ([TNFRSF8 Products](#))

Molecular Weight: 105-120kDa

Gene ID: 943

UniProt: [P28908](#)

## Application Details

---

Application Notes: Positive Control: Hodgkin's lymphoma.  
Known Application: Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

---

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months



#### Immunohistochemistry

**Image 1.** Formalin-fixed, paraffin-embedded human Hodgkin's lymphoma stained with CD30 Ab (CD30/412).



#### Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Hodgkin's lymphoma stained with CD30 Ab (CD30/412).