

Datasheet for ABIN6941166

**Recombinant anti-NAPSA antibody (AA 189-299)****2** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	NAPSA
Binding Specificity:	AA 189-299
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This NAPSA antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant human Napsin-A protein fragment (aa189-299) (exact sequence is proprietary)
Clone:	NAPSA-1865R
Isotype:	IgG
Purification:	Purified by Protein A/G

## Target Details

Target:	NAPSA
Alternative Name:	NAPSA ( <a href="#">NAPSA Products</a> )
Background:	Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein

## Target Details

B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high expression levels of napsin A in human lung and kidney but low expression in spleen. Napsin A is expressed in type II pneumocytes and in adenocarcinomas of lung. The high specificity expression of napsin A in adenocarcinomas of lung is useful to distinguish primary lung adenocarcinomas from adenocarcinomas of other organs.

Molecular Weight: 37kDa

Gene ID: 9476

UniProt: [O96009](#)

Pathways: [Tube Formation](#), [Asymmetric Protein Localization](#), [Embryonic Body Morphogenesis](#)

## Application Details

Application Notes: Positive Control: Lung adenocarcinoma.  
Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months



### SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified Napsin A Rabbit Recombinant Monoclonal Antibody (NAPSA/1865R).



### Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Lung Adenocarcinoma stained with Napsin A Rabbit Recombinant Monoclonal Antibody (NAPSA/1865R).