

Datasheet for ABIN6941220
anti-CD59 antibody



[Go to Product page](#)

2 Images

Overview

| | |
|--------------|---|
| Quantity: | 100 µg |
| Target: | CD59 |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This CD59 antibody is un-conjugated |
| Application: | Flow Cytometry (FACS), Immunofluorescence (IF), Functional Studies (Func) |

Product Details

| | |
|--------------|---|
| Immunogen: | Stimulated human leukocytes |
| Clone: | 193-27 |
| Isotype: | IgM kappa |
| Specificity: | <p>Reacts with human CD59, a 20 kDa glycosyl phosphatidyl-inositol (GPI)-anchored cell surface protein (Workshop VI, Code N-L036). CD59 regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. CD59 is widely distributed on cells in all tissues. It inhibits formation of MAC, thus protecting cells from complement-mediated lysis. The expression of CD59 on erythrocytes is important for their survival. Genetic defects in GPI-anchor attachment, that cause a reduction or loss of CD59 and CD55 on erythrocytes produce the symptoms of the disease paroxysmal hemoglobinuria (PNH). This MAbs recognizes CD59</p> |

Product Details

transfected cells. It is useful for study on GPI-anchored proteins, PNH and CD59 functions.

No Cross-Reactivity: Baboon, Horse (Equine)

Target Details

Target: CD59

Alternative Name: CD59 ([CD59 Products](#))

Molecular Weight: 20kDa

Gene ID: 966

UniProt: [P13987](#)

Pathways: [Complement System](#)

Application Details

Application Notes: Positive Control: Jurkat or Raji cells. Human lymphocytes. Human lymph node and tonsil.
Known Application: Functional Studies (Order Ab without Azide),Flow Cytometry (0.5-1 µg/million cells), ,Immunofluorescence (0.5-1 µg/mL), ,Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10mM PBS with 0.05% BSA & 0.05% azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months



Immunohistochemistry

Image 1. Formalin-fixed, paraffin-embedded human bladder stained with CD59 Mouse Monoclonal Antibody (193-27).



Immunohistochemistry

Image 2. Formalin-fixed, paraffin-embedded human bladder stained with CD59 Mouse Monoclonal Antibody (193-27).