

Datasheet for ABIN6941289

**anti-TNFSF15 antibody**

3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	TNFSF15
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNFSF15 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Staining Methods (StM)

## Product Details

Immunogen:	Recombinant human VEGI fragment
Clone:	VEGI-1283
Isotype:	IgG1 kappa
Purification:	Purified by Protein A/G

## Target Details

Target:	TNFSF15
Alternative Name:	TNFSF15 ( <a href="#">TNFSF15 Products</a> )
Background:	VEGI is an anti-angiogenic cytokine that belongs to tumor necrosis factor superfamily, member 15 (TNFSF15). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate

## Target Details

NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation, and thus may function as an angiogenesis inhibitor. Reduced expression of VEGF has been reported as a marker of poor prognosis in breast cancer.

Molecular Weight: 22kDa

Gene ID: 9966

UniProt: [O95150](#)

Pathways: [Positive Regulation of Endopeptidase Activity](#), [Autophagy](#)

## Application Details

Application Notes: Positive Control: Endothelial cells. Colon, intestine, placenta, lung, liver, kidney, pancreas, spleen and prostate.

Known Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes) Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

## Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -80 °C


Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months




### SDS-PAGE

**Image 1.** SDS-PAGE Analysis Purified TNFS15 / VEGI Monoclonal Antibody (VEGI /1283). Confirmation of Purity and Integrity of Antibody



### Immunohistochemistry

**Image 2.** Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with TNFS15 / VEGI Monoclonal Antibody (VEGI /1283).



### Immunohistochemistry

**Image 3.** Formalin-fixed, paraffin-embedded human Parathyroid Mass stained with TNFS15 / VEGI Monoclonal Antibody (VEGI /1283).