

Datasheet for ABIN6941298

Recombinant anti-E-cadherin antibody

5 Images

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	E-cadherin (CDH1)
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This E-cadherin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Staining Methods (StM), Immunostaining (Ist)

Product Details

Immunogen:	Recombinant full-length human E-Cadherin protein
Clone:	RCDH1-1525
Isotype:	IgG1 kappa
Specificity:	Recognizes a protein of 120-80 kDa, identified as E-cadherin. Cadherins comprise a family of Ca ²⁺ -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH ₂ terminal repeats. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as -catenin, to regulate cadherin function. E-cadherin plays an important role in epithelial cell adhesion. A decreased expression of E-cadherin is associated

Product Details

with metastatic potential and poor prognosis in breast cancer, prostate and esophageal cancer. In combination with p120 Catenin, it is useful for the differentiation between ductal (E-cadherin +) and lobular (E-cadherin -) breast carcinomas. It may also help in diagnosis of mesothelioma.

No Cross-Reactivity: Mouse (Murine), Rat (Rattus)

Purification: Purified by Protein A/G

Target Details

Target: E-cadherin (CDH1)

Alternative Name: CDH1 ([CDH1 Products](#))

Molecular Weight: 120-80kDa (Mature), 135kDa (Precursor)

Gene ID: 999

UniProt: [P12830](#)

Pathways: [WNT Signaling](#), [Sensory Perception of Sound](#), [Cell-Cell Junction Organization](#), [Tube Formation](#)

Application Details

Application Notes: Positive Control: MCF-7, LS174T, Raji, HT29, SK-BR3 cells. Prostate or Colon carcinomas.
Known Application: Western Blot (0.5-1 µg/mL), Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 10-20 min followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C,-80 °C

Handling

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


Expiry Date: 24 months

Images




Western Blotting

Image 1. Western Blot Analysis of MCF-7 cell lysate using E-Cadherin Mouse Recombinant Monoclonal Antibody (rCDH1/1525).



Protein Array

Image 2. Analysis of Protein Array containing >19,000 full-length human proteins using E-Cadherin Mouse Monoclonal Antibody (rCDH1/1525) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Immunohistochemistry

Image 3. Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with E-Cadherin Mouse Recombinant Monoclonal Antibody (rCDH1/1525).

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN6941298.