

Datasheet for ABIN6941366
anti-BrdU antibody



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	BrdU
Reactivity:	Please inquire
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This BrdU antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Flow Cytometry (FACS), Immunofluorescence (IF), Neutralization (Neut), Staining Methods (StM)

Product Details

Immunogen:	Bromodeoxyuridine (BrdU) conjugated to KLH
Clone:	SPM166
Isotype:	IgG1 kappa
Specificity:	It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting S-phase cells, providing useful information on the aggressiveness of tumors.
Purification:	Purified by Protein A/G

Target Details

Target: BrdU

Abstract: [BrdU Products](#)

Target Type: Chemical

Molecular Weight: 307Da

Application Details

Application Notes: Positive Control: Cells grown in presence of BrdU or tissues from experimental animals injected with BrdU.

Known Application: Flow Cytometry (0.5-1 µg/million cells), Immunofluorescence (0.5-1 µg/mL), Immunohistochemistry (Formalin-fixed) (0.5-1 µg/mL for 30 minutes at RT) (Use Ab at 1 µg/mL for 30 minute at RT)[For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion with trypsin at 1 mg/mL PBS, 10 min at 37oC
Optimal dilution for a specific application should be determined.

Restrictions: For Research Use only

Handling

Concentration: 200 µg/mL

Buffer: 10 mM PBS with 0.05 % BSA & 0.05 % azide.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C,-80 °C

Storage Comment: Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Expiry Date: 24 months



Immunohistochemistry

Image 1. Formalin-fixed, paraffin-embedded Mouse Intestine stained with BrdU Monoclonal Antibody (SPM166).