

## Datasheet for ABIN6941397 **anti-CD176 antibody**



[Go to Product page](#)

### Overview

Quantity:	100 µg
Target:	CD176
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD176 antibody is un-conjugated
Application:	ELISA, Coating (Coat)

### Product Details

Immunogen:	Neuraminidase-treated human red blood cells
Clone:	A68-B-A11
Isotype:	IgM kappa
Specificity:	Recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivity with sialylated glycophorin. The Thomsen-Friedenreich antigen acts as an oncofetal antigen, with low expression in normal adult tissues but increasing to fetal levels of expression in hyperplasia or malignancy. It is considered as a pan-carcinoma marker. This MAb is capable to agglutinate desialylated red blood cells. During metastasis, the ability of malignant cells to form multicellular aggregates via homotypic or heterotypic aggregation and their adhesion to the endothelium are critical. The tumor-associated carbohydrate Thomsen-Friedenreich antigen (Gal-GalNAc) is involved in tumor cell adhesion and tissue invasion. It also causes an immune response, and overexpression of the

## Product Details

antigen causes cancer cells to be more sensitive to natural killer cell lysis. The Thomsen-Friedenreich antigen is suppressed in normal healthy cells and represents one of the few chemically well-defined antigens associated with tumor malignancy. The presence of the Thomsen-Friedenreich antigen on the surface of cancer cells may result from a divergence from the normal pathway for O-linked glycosylation in these cells, most likely caused by inappropriate localization of the enzymes involved in synthesis of the disaccharide.

## Target Details

Target:	CD176
Alternative Name:	Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker) ( <a href="#">CD176 Products</a> )

## Application Details

Application Notes:	Positive Control: KG1 cells. Human colorectal carcinoma tissues. Known Application: ELISA (For coating use Ab at 1-5 µg/mL, order Ab without BSA), Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only

## Handling

Concentration:	200 µg/mL
Buffer:	10 mM PBS with 0.05 % BSA & 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Expiry Date:	24 months