antibodies -online.com







Overview

Quantity:	100 μL
Target:	ADAP2
Binding Specificity:	AA 201-300
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADAP2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Centaurin alpha 2
Isotype:	IgG
Specificity:	Possible cross-reactivity with ADAP1 (85 % homology with the immunogen sequence, 43 kDa)
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ADAP2
Alternative Name:	Centaurin alpha 2 (ADAP2 Products)
Background:	Synonyms: ADAP2, ADAP2_HUMAN, Arf-GAP with dual PH domain-containing protein 2, ArfGAF
	with dual PH domains 2, cent b, Centa 2, CENTA2, Centaurin alpha 2, Centaurin alpha 2 protein,
	Centaurin beta, Centaurin-alpha-2, Cnt-a2, HSA272195.
	Background: The ADP-ribosylation factor (ARF) family of small GTP-binding proteins are
	involved in vesicular transport regulation and in controlling cytoskeletal organization and cell
	adhesion. The Centaurin GTPase-activating protein family comprise a subset of ARF regulatory
	molecules that transduce PI 3-kinase activation into coordinated control of ARF-dependent
	pathways. This family includes ASAP1, ACAP1, ACAP2, AGAP1, ARAP1, ARAP2, Centaurin a1,
	Centaurin g3 and the recently discovered Centaurin a2. Expressed in a wide variety of tissues
	such as fat, heart and skeletal muscle, Centaurin a2 is thought to negatively regulate ARF-
	mediated actin rearrangement by binding activated PI 3-kinase. Although the exact function of
	Centaurin a2 is not yet known, its high sequence similarity with Centaurin a1 suggests that it
	may also act as an ARF6 GTPase.
Gene ID:	55803
UniProt:	Q9NPF8
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months