

[Go to Product page](#)

Datasheet for ABIN6942506
anti-ADAP2 antibody (AA 201-300) (Biotin)

Overview

Quantity:	100 µL
Target:	ADAP2
Binding Specificity:	AA 201-300
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADAP2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Centaurin alpha 2
Isotype:	IgG
Specificity:	Possible cross-reactivity with ADAP1 (85 % homology with the immunogen sequence, 43 kDa)
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ADAP2
---------	-------

Target Details

Alternative Name:	Centaurin alpha 2 (ADAP2 Products)
Background:	<p>Synonyms: ADAP2, ADAP2_HUMAN, Arf-GAP with dual PH domain-containing protein 2, ArfGAP with dual PH domains 2, cent b, Centa 2, CENTA2, Centaurin alpha 2, Centaurin alpha 2 protein, Centaurin beta, Centaurin-alpha-2, Cnt-a2, HSA272195.</p> <p>Background: The ADP-ribosylation factor (ARF) family of small GTP-binding proteins are involved in vesicular transport regulation and in controlling cytoskeletal organization and cell adhesion. The Centaurin GTPase-activating protein family comprise a subset of ARF regulatory molecules that transduce PI 3-kinase activation into coordinated control of ARF-dependent pathways. This family includes ASAP1, ACAP1, ACAP2, AGAP1, ARAP1, ARAP2, Centaurin a1, Centaurin g3 and the recently discovered Centaurin a2. Expressed in a wide variety of tissues such as fat, heart and skeletal muscle, Centaurin a2 is thought to negatively regulate ARF-mediated actin rearrangement by binding activated PI 3-kinase. Although the exact function of Centaurin a2 is not yet known, its high sequence similarity with Centaurin a1 suggests that it may also act as an ARF6 GTPase.</p>
Gene ID:	55803
UniProt:	Q9NPF8

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months