

[Go to Product page](#)

Datasheet for ABIN6942509

anti-Centaurin beta 2 antibody (AA 101-200) (Biotin)

Overview

Quantity:	100 µL
Target:	Centaurin beta 2 (ACAP2)
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Centaurin beta 2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Centaurin beta 2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Centaurin beta 2 (ACAP2)
Alternative Name:	Centaurin beta 2 (ACAP2 Products)
Background:	Synonyms: ACAP 2, ACAP2, ACAP2_HUMAN, ANK repeat and PH domain-containing protein 2,

Target Details

Ankyrin repeat, Arf GAP with coiled coil ANK repeat and PH domains 2, ARF GAP with coiled coil, Arf-GAP with coiled-coil, CENT B 2, Centaurin beta2, Centaurin-beta-2, CENTB 2, CENTB2, Cnt b2, Cnt-b2, Cntb2, KIAA0041, Pleckstrin homology domains 2, Similar to yeast ZINC FINGER PROTEIN GCS1.

Background: The ADP-ribosylation factor (ARF) family of small GTP-binding proteins are involved in vesicular transport regulation and in controlling cytoskeletal organization and cell adhesion. These proteins mainly regulate membrane traffic. ACAP2 is a member of the centaurin GTPase-activating protein (GAP) family, which comprises a subset of ARF regulatory molecules that transduce PI 3-kinase activation into coordinated control of ARF-dependent pathways. ACAP1 and ACAP2 are both widely expressed in peripheral, tubular membranes and usually interact with each other in various tissues. GAP activity of both ACAP1 and ACAP2 is dependent upon phosphatidylinositol 4,5-bisphosphate [PtdIns(4,5)P₂]. ACAP2 associates with ARF1 and ARF6. Overexpression of ACAP2 blocks the formation of ARF6-dependent protrusions. K1L is a protein required for growth of the Vaccinia Virus that interacts with the ankyrin repeats of ACAP2.

Gene ID: 23527

UniProt: [Q15057](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months