# antibodies -online.com





## anti-CES3 antibody (AA 431-530)



$\sim$			
	N/6	1//r	$I \cap V$

Quantity:	100 μL
Target:	CES3
Binding Specificity:	AA 431-530
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CES3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CES3
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

### Target Details

Target:	CES3
Alternative Name:	CES3 (CES3 Products)

### Target Details

Background:	Synonyms: Carboxylesterase 3, CES 3, Ces3, ES3, EST3_HUMAN, FLJ21736, Liver	
	carboxylesterase 31 homolog.	
	Background: This gene encodes a member of the carboxylesterase large family. The family	
	members are responsible for the hydrolysis or transesterification of various xenobiotics, such	
	as cocaine and heroin, and endogenous substrates with ester, thioester, or amide bonds. They	
	may participate in fatty acyl and cholesterol ester metabolism, and may play a role in the blood-	
	brain barrier system. This gene is expressed in several tissues, particularly in colon, trachea and	
	in brain, and the protein participates in colon and neural drug metabolism. Multiple alternatively	
	spliced transcript variants encoding distinct isoforms have been reported, but the biological	
	validity and/or full-length nature of some variants have not been determined.[provided by	
	RefSeq, Jun 2010]	
Gene ID:	23491	
UniProt:	Q6UWW8	
Pathways:	Monocarboxylic Acid Catabolic Process	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

### Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months