# antibodies -online.com





## anti-CYP27B1 antibody (AA 1-100) (Biotin)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	CYP27B1
Binding Specificity:	AA 1-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP27B1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CYP27B1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	CYP27B1	
Alternative Name:	CYP27B1 (CYP27B1 Products)	
Background:	Synonyms: 1alpha(OH)ase, 25-hydroxyvitamin D(3) 1-alpha-hydroxylase, 25-hydroxyvitamin D-1	

alpha hydroxylase, 25-OHD-1 alpha-hydroxylase, Calcidiol 1-monooxygenase, CP27B\_HUMAN, CP2B, CYP1, CYP1ALPHA, CYP27B, Cyp27b1, Cytochrome p450 27B1, Cytochrome p450 27B13, Cytochrome P450 family 27 subfamily B polypeptide 1, Cytochrome P450 subfamily XXVIIB (25-hydroxyvitamin D-1-alpha-hydroxylase) polypeptide 1, Cytochrome P450 subfamily XXVIIB polypeptide 1, Cytochrome P450C1 alpha, Cytochrome P450VD1-alpha, mitochondrial, P450C1 alpha, P450C1, P450C1-alpha, P450VD1-alpha, PDDR, VD3 1A hydroxylase, VDD1, VDDR, VDDR I, VDDRI, VDR.

Background: The cytochrome P450 proteins are monooxygenases that catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. P450 enzymes are classified into subfamilies based on their sequence similarities. CYP27B1, a 508-amino acid protein that belongs to the XXVIIB subfamily of the cytochrome P450 family, localizes to the mitochondrion and is expressed in the kidney. The CYP27B1 protein catalyzes the conversion of 25-hydroxyvitamin D3 (25(OH)D) to 1-alpha,25-dihydroxyvitamin D3 (1,25(OH)2D) and functions in calcium metabolism, normal bone growth, and tissue differentiation. Mutations in the gene which encodes for CYP27B1 cause vitamin D-dependent rickets type 1 (VDDR-1), also designated pseudovitamin D deficiency rickets (PDDR), an autosomal recessive disease characterized by early onset of rickets with hypocalcemia and muscle weakness.

Gene ID: 1594

UniProt: 015528

Pathways: Metabolism of Steroid Hormones and Vitamin D

IHC-P 1:200-400

#### **Application Details**

Application Notes: WB 1:300-5000

Restrictions: For Research Use only

#### Handling

Format:

Concentration:

1 μg/μL

Buffer:

Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

## Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months