# antibodies -online.com





# anti-CYP2C19 antibody (AA 201-300)



Go to Product page

( )	11/0	K\ /	iew	1
	$\cup$	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	CYP2C19
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP2C19 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CYP2C19	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	CYP2C19
Alternative Name:	CYP2C19 (CYP2C19 Products)

#### **Target Details**

Background:

Synonyms: Al266900, CYP2C, CYP2C19, CYP2C37, CYP2C50, Cyp2c54, CYP450-2C, CPCJ, CYPIIC17, CYPIIC19, EG404195, LOC293989, MGC109053, P450-11A, P450C2C, P450IIC19, CP2CJ\_HUMAN Cytochrome P450 2C19, R)-limonene 6-monooxygenase, (S)-limonene 6-monooxygenase, (S)-limonene 7-monooxygenase, CYPIIC17, CYPIIC19, Cytochrome P450-11A, Cytochrome P450-254C, Mephenytoin 4-hydroxylase.

Background: This gene encodes a member of the cytochrome P450 superfamily of enzymes and is responsible for the metabolism of a number of therapeutic agents such as the anticonvulsant drug S-mephenytoin, omeprazole, proguanil, certain barbiturates, diazepam, propranolol, citalopram, and imipramine. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, omeprazole, diazepam and some barbiturates. Polymorphism within this gene is associated with variable ability to metabolize mephenytoin, known as the poor metabolizer and extensive metabolizer phenotypes. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. [provided by RefSeq]

Gene ID:

1557

UniProt:

P33261

#### **Application Details**

**Application Notes:** 

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

#### Handling

Format:

Liquid

Concentration:

 $1 \mu g/\mu L$ 

## Handling

Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	