Datasheet for ABIN6942779

anti-CYP2C19 antibody (AA 201-300) (AbBy Fluor® 594)

Go to Product page

Overview

Quantity:	100 µL
Target:	CYP2C19
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP2C19 antibody is conjugated to AbBy Fluor® 594
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CYP2C19
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	CYP2C19
Alternative Name:	CYP2C19 (CYP2C19 Products)
Background:	Synonyms: AI266900, CYP2C, CYP2C19, CYP2C37, CYP2C50, Cyp2c54, CYP450-2C, CPCJ,

Target Details

	CYPIIC17, CYPIIC19, EG404195, LOC293989, MGC109053, P450-11A, P450C2C, P450IIC19, CP2CJ_HUMAN Cytochrome P450 2C19, R)-limonene 6-monooxygenase, (S)-limonene 6-monooxygenase, (S)-limonene 7-monooxygenase, CYPIIC17, CYPIIC19, Cytochrome P450-11A, Cytochrome P450-254C, Mephenytoin 4-hydroxylase. Background: This gene encodes a member of the cytochrome P450 superfamily of enzymes and is responsible for the metabolism of a number of therapeutic agents such as the anticonvulsant drug S-mephenytoin, omeprazole, proguanil, certain barbiturates, diazepam, propranolol, citalopram, and imipramine. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, omeprazole, diazepam and some barbiturates. Polymorphism within this gene is associated with variable ability to metabolize mephenytoin, known as the poor metabolizer and extensive metabolizer phenotypes. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. [provided by RefSeq]
Gene ID:	1557
UniProt:	P33261

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200

Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Copyright antibodies-online. All rights reserved.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

 Copyright antibodies-online. All rights reserved.