antibodies .- online.com





anti-DHX30 antibody (AA 1-100) (Alexa Fluor 488)



| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | DHX30 | |
| Binding Specificity: | AA 1-100 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This DHX30 antibody is conjugated to Alexa Fluor 488 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human DHX30 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | DHX30 |
|-------------------|------------------------|
| Alternative Name: | DHX30 (DHX30 Products) |

Target Details

| Background: | Synonyms: DDX30, DEAH box protein 30, Dhx30, DHX30_HUMAN, FLJ11214, KIAA0890, | |
|---------------------|---|--|
| | Putative ATP-dependent RNA helicase DHX30, Ret CoR. | |
| | Background: DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), | |
| | are putative RNA helicases. They are implicated in a number of cellular processes involving | |
| | alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial | |
| | splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some | |
| | members of this DEAD box protein family are believed to be involved in embryogenesis, | |
| | spermatogenesis, and cellular growth and division. The family member encoded by this gene is | |
| | a mitochondrial nucleoid protein that associates with mitochondrial DNA. It has also been | |
| | identified as a component of a transcriptional repressor complex that functions in retinal | |
| | development, and it is required to optimize the function of the zinc-finger antiviral protein. | |
| | Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Feb | |
| | 2013] | |
| Gene ID: | 22907 | |
| UniProt: | Q7L2E3 | |
| Pathways: | Chromatin Binding | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| Precaution of Use: | | |
| Precaution of Use: | handled by trained staff only. | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |