

### Datasheet for ABIN6943006

# anti-DLG4 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL
Target:	DLG4
Reactivity:	Rat, Mouse, Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This DLG4 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

### **Product Details**

Immunogen:	Recombinant human PSD95 around N-terminus		
Clone:	1H8		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse, Rat		
Purification:	Purified by Protein A.		
Toward Dataila			

#### Target Details

Target:	DLG4
Alternative Name:	DLG4 (DLG4 Products)

## Target Details

-			
Background:	Synonyms: Disks large homolog 4, Postsynaptic density protein 95, Synapse-associated protein		
	90, PSD-95, SAP-90, SAP90, Disks large homolog 4, discs large MAGUK scaffold protein 4, DLG4		
	Background: Interacts with the cytoplasmic tail of NMDA receptor subunits and shaker-type		
	potassium channels. Required for synaptic plasticity associated with NMDA receptor signaling.		
	Overexpression or depletion of DLG4 changes the ratio of excitatory to inhibitory synapses in		
	hippocampal neurons. May reduce the amplitude of ASIC3 acid-evoked currents by retaining		
	the channel intracellularly. May regulate the intracellular trafficking of ADR1B (By similarity).		
Gene ID:	1742		
UniProt:	P78352		
Pathways:	Regulation of Muscle Cell Differentiation, Synaptic Membrane, Skeletal Muscle Fiber		
	Development, Asymmetric Protein Localization, Regulation of long-term Neuronal Synaptic		
	Plasticity		
Application Details			
Application Notes:	WB 1:300-5000		
	FCM 1:20-100		
	IHC-P 1:200-400		
	IF(IHC-P) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 1xTBS (pH 7.4), 1 % BSA, 40 %Glycerol and 0.05 %		
	Sodium Azide.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		

	1	l:
-	ココロロ	urnen
	Hand	III IU

Expiry Date:

12 months