# antibodies - online.com





Datasheet for ABIN6943027

## anti-DNAJC19 antibody (AA 21-116)



( )	ve	K\ /		A .
	$\cup$	1 V/	-	V۷

Quantity:	100 μL	
Target:	DNAJC19	
Binding Specificity:	AA 21-116	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DNAJC19 antibody is un-conjugated	
Application:	eplication: ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozer Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DNAJC19
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

|--|

### Target Details

Alternative Name:	DNAJC19 (DNAJC19 Products)	
Background:	Synonyms: DnaJ (Hsp40) homolog, subfamily C, member 19, DnaJ homolog subfamily C	
	member 19, DNAJC19, homolog of yeast TIM14, Mitochondrial import inner membrane	
	translocase subunit TIM14, PAM18, TIM 14, TIM14, TIM14_HUMAN, TIMM 14, TIMM14,	
	translocase of the inner mitochondrial membrane 14, Translocase of the inner mitochondrial	
	membrane 14, yeast homolog of.	
	Background: The protein encoded by this gene is thought to be part of a complex involved in the	
	ATP-dependent transport of transit peptide-containing proteins from the inner cell membrane	
	to the mitochondrial matrix. Defects in this gene are a cause of 3-methylglutaconic aciduria	
	type 5 (MGA5), also known as dilated cardiomyopathy with ataxia (DCMA). Alternative splicing	
	of this gene results in multiple transcript variants. Related pseudogenes have been identified on	
	chromosomes 1, 2, 6, 10, 14 and 19. [provided by RefSeq, Jan 2012]	
Gene ID:	131118	
UniProt:	Q96DA6	
Pathways:	SARS-CoV-2 Protein Interactome	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

### Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months