# antibodies -online.com





# anti-EN1 antibody (AA 351-392) (Alexa Fluor 750)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	EN1	
Binding Specificity:	AA 351-392	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EN1 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human EN1/Engrailed 1	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	EN1	
Alternative Name:	EN1/Engrailed 1 (EN1 Products)	

#### **Target Details**

Background:

Synonyms: En 1, EN1, Engrailed 1, Engrailed, Engrailed homeobox 1, Engrailed homolog 1, HME1\_HUMAN, Homeobox protein en-1, Homeobox protein engrailed 1, Homeobox protein engrailed-1, Hu-En-1, Mo-en-1.

Background: The engrailed-1 gene, EN1, a murine homolog of the Drosophila homeobox gene engrailed (EN), is required for midbrain and cerebellum development and dorsal/ventral patterning of the limbs as well as apical ectodermal ridge formation. In Drosophila, the EN gene plays an important role during development in segmentation, where it is required for the formation of posterior compartments. Human EN-1 and EN-2 are homeodomain-containing proteins and have been implicated in the control of pattern formation during development of the central nervous system. Different mutations in the mouse homo-logs, EN-1 and EN-2, produce different developmental defects that frequently are lethal. EN-1 is highly expressed by essentially all dopaminergic neurons in the substantia nigra and ventral tegmentum. EN-1 and EN-2 regulate expres-sion of a-synuclein, a gene that is genetically linked to Parkinson?s disease.

Gene ID:

2019

UniProt:

Q05925

Pathways:

Dopaminergic Neurogenesis

#### **Application Details**

Application Notes:

FCM 1:20-100

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	

Liquid

Concentration:

 $1 \mu g/\mu L$ 

Buffer:

Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

50 % Glycerol.

Preservative:

ProClin

Precaution of Use:

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

## Handling

	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	