

Datasheet for ABIN6943328

anti-EXOC1 antibody (AA 701-800) (HRP)



Go to Product page

| (| ۱۱/ | e | r\/ | Ì١ | | ۱۸ | |
|---|-----|--------|-----|----|---------------|----|---|
| | , v | \cup | V | 1 | $\overline{}$ | V | V |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | EXOC1 |
| Binding Specificity: | AA 701-800 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EXOC1 antibody is conjugated to HRP |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human EXOC1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Cow,Sheep,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | EXOC1 |
|-------------------|------------------------|
| Alternative Name: | EXOC1 (EXOC1 Products) |

Target Details

| Background: | Synonyms: BM-012, exoc1, EXOC1_HUMAN, Exocyst complex component 1, Exocyst complex |
|---------------------|--|
| | component Sec3, FLJ10893, SEC3, SEC3 like 1, SEC3L1, SEC3P. |
| | Background: Exocytosis is crucial in membrane trafficking and it mediates hormone and |
| | neurotransmitter secretion out of the cell, as well as the incorporation of membrane proteins |
| | and lipids to the plasma membrane. It is crucial for cell-cell communication, cell growth and cell |
| | polarity. The exocyst complex is a multi-protein complex that consists of Sec3, Sec5, Sec6, |
| | Sec8, Sec10, Sec15, Exo70 and Exo84, and is essential for targeting exocytic vesicles to |
| | specific docking sites on the plasma membrane. Exo70, also known as EXOC7 (exocyst |
| | complex component 7), EXOC1 or 2-5-3p, is a 735 amino acid peripheral membrane protein that |
| | is a component of the exocyst complex. Localized to the cytoplasm and the cell membrane, |
| | Exo70 plays an essential role in the docking of exocystic vesicles to target sites on the plasma |
| | membrane and, specifically, may be involved in insulin-induced protein docking within the cell. |
| | Four isoforms of Exo70 are expressed due to alternative splicing events. |
| Gene ID: | 55763 |
| UniProt: | Q9NV70 |
| Pathways: | Peptide Hormone Metabolism, Synaptic Vesicle Exocytosis |
| Application Details | |
| Application Notes: | IHC-P 1:200-400 |
| | IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |