antibodies -online.com





anti-EXOC1 antibody (AA 701-800) (Alexa Fluor 350)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	EXOC1	
Binding Specificity:	AA 701-800	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EXOC1 antibody is conjugated to Alexa Fluor 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EXOC1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	EXOC1
Alternative Name:	EXOC1 (EXOC1 Products)

Target Details

Synonyms: BM-012, exoc1, EXOC1_HUMAN, Exocyst complex component 1, Exocyst complex
component Sec3, FLJ10893, SEC3, SEC3 like 1, SEC3L1, SEC3P.
Background: Exocytosis is crucial in membrane trafficking and it mediates hormone and
neurotransmitter secretion out of the cell, as well as the incorporation of membrane proteins
and lipids to the plasma membrane. It is crucial for cell-cell communication, cell growth and cell
polarity. The exocyst complex is a multi-protein complex that consists of Sec3, Sec5, Sec6,
Sec8, Sec10, Sec15, Exo70 and Exo84, and is essential for targeting exocytic vesicles to
specific docking sites on the plasma membrane. Exo70, also known as EXOC7 (exocyst
complex component 7), EXOC1 or 2-5-3p, is a 735 amino acid peripheral membrane protein that
is a component of the exocyst complex. Localized to the cytoplasm and the cell membrane,
Exo70 plays an essential role in the docking of exocystic vesicles to target sites on the plasma
membrane and, specifically, may be involved in insulin-induced protein docking within the cell.
Four isoforms of Exo70 are expressed due to alternative splicing events.
55763
Q9NV70
Peptide Hormone Metabolism, Synaptic Vesicle Exocytosis
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months