antibodies -online.com





anti-EXOC3L2 antibody (AA 151-250)



_	
()Ver	view

Quantity:	100 μL
Target:	EXOC3L2
Binding Specificity:	AA 151-250
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOC3L2 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EXOC3L2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow
Purification:	Purified by Protein A.

Target Details

rarget. EXOCSE2

Target Details

Alternative Name:	EXOC3L2 (EXOC3L2 Products)
Background:	Synonyms: Exocyst complex component 3 like, Exocyst complex component 3 like protein 2,
	FLJ36147, EX3L2_HUMAN, HBV X transactivated gene 7 protein, HBV XAg transactivated
	protein 7, MGC16332, OTTHUMP00000207128, OTTMUSP00000045827, Protein 7
	transactivated by hepatitis B virus X antigen (HBxAg), XTP7.
	Background: Exocytosis is crucial in membrane trafficking and it mediates hormone and
	neurotransmitter secretion out of the cell, as well as the incorporation of membrane proteins
	and lipids to the plasma membrane. It is crucial for cell-cell communication, cell growth and ce
	polarity. The exocyst complex is a multi-protein complex that consists of Sec3, Sec5, Sec6,
	Sec8, Sec10, Sec15, Exo70 and Exo84, and is essential for targeting exocytic vesicles to
	specific docking sites on the plasma membrane. The exocyst complex inhibits tubulin
	polymerization in vitro, suggesting that the exocyst complex is important for modulating the
	microtubule dynamics that underlie exocytosis. ExoC3L2 (Exocyst complex component 3-like
	protein 2), also known as HBV X-transactivated gene 7 protein, is a 409 amino acid protein that
	is related to the Sec6 component of the exocyst complex.
Gene ID:	90332
UniProt:	Q2M3D2
Application Details	
Application Notes:	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months