# antibodies -online.com





## anti-EXOSC4 antibody (AA 51-150) (Biotin)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL
Target:	EXOSC4
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC4 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EXOSC4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	EXOSC4
Alternative Name:	EXOSC4 (EXOSC4 Products)

#### Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		$\sim$	<i>.</i>	ч.

Synonyms: EXOSC 4, Exosome complex component RRP41, Exosome complex exonuclease RRP41, Exosome component 4, Exosome component Rrp41, FLJ20591, hRrp41p, p12A, EXOS4\_HUMAN, Ribosomal RNA processing protein 41, RRP 41, RRP41A, Rrp41A, Rrp41p, SKI 6, SKI6, Ski6p.

Background: The exosome is a multisubunit complex composed of several highly conserved subunits, some of which are 3 to 5 exoribonucleases. The complex is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich (ARE) elements in their untranslated 3 region. EXOSC4 (exosome component 4), also known as SKI6, RRP41 or p12A, is a 245 amino acid protein that localizes to both the nucleus and the cytoplasm and shares 96 % sequence identity with its mouse counterpart. Functioning as a component of the exosome complex, EXOSC4 exhibits 3'-5' exonuclease activity and is required for the 3'-processing of 7S pre-RNA to mature 5.8S rRNA. The gene encoding EXOSC4 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.

Gene ID:

54512

UniProt:

Q9NPD3

#### **Application Details**

Application Notes:

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months