## antibodies -online.com





Background:

## anti-FPA antibody (AA 20-120) (Alexa Fluor 350)



Overview		
Quantity:	100 μL	
Target:	FPA	
Binding Specificity:	AA 20-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FPA antibody is conjugated to Alexa Fluor 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Fibrinopeptide A	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	
Target Details		
Target:	FPA	
Alternative Name:	Fibrinopeptide A (FPA Products)	

Synonyms: FGA, Fib2, FIBA\_HUMAN, Fibrinogen alpha chain, fibrinogen alpha chain isoform

alpha preproprotein, Fibrinogen alpha/alpha E chain [Precursor], fibrinogen alpha chain isoform
alpha-E preproprotein, fibrinogen alpha chain isoform alpha preproprotein.

Background: The protein encoded by this gene is the alpha component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia, afibrinogenemia and renal amyloidosis. Alternative splicing results in two isoforms which vary in the carboxy-terminus. [provided by RefSeq, Jul 2008]

Gene ID: 2243

UniProt: P02671

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	