# antibodies - online.com







# anti-IQSEC1 antibody (AA 451-550)



$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	IQSEC1
Binding Specificity:	AA 451-550
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IQSEC1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GEP100
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target: IQSEC1
----------------

### Target Details

rarget betails	
Alternative Name:	GEP100/IQSEC1 (IQSEC1 Products)
Background:	Synonyms: ADP-ribosylation factors guanine nucleotide-exchange protein 100, ADP-ribosylation
	factors guanine nucleotide-exchange protein 2, ARF GEP100, ARFGEP100, BRAG2, Brefeldin A
	resistant ARF GEF2, Brefeldin-resistant Arf-GEF 2 protein, GEP100, IQ motif and Sec7 domain 1
	IQ motif and SEC7 domain-containing protein 1, IQEC1_HUMAN, Iqsec1, Kiaa0763.
	Background: The ADP-ribosylation factor (ARF) protein family are structurally and functionally
	conserved members of the Ras superfamily of regulatory GTP-binding proteins. ARFs influence
	vesicle trafficking and signal transduction in eukaryotic cells. ARF6 plays a role in protein
	trafficking near the plasma membrane, including receptor recycling, cell adhesion and cell
	migration. ARF6 colocalizes with the ARF guanine nucleotide-exchange protein (GEP) BRAG2,
	also designated GEP100. BRAG2 is ubiquitously expressed as two isoforms, BRAG2a and
	BRAG2b, which can cycle between the cytoplasm and the nucleus. BRAG2, via its interaction
	with ARF6, is involved in the regulation of cell adhesion by controlling Integrin 1 endocytosis
	and E-cadherin redistribution. BRAG2 has also been shown to bind directly to Tyr1068/1086-
	phosphorylated EGFR to activate ARF6, which induces tumor invasion in MCF7 cells. Therefore
	BRAG2 may contribute to the metastasis and malignancy of some breast cancer cells.
Gene ID:	9922
UniProt:	Q6DN90
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL

## Handling

Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	