# antibodies -online.com





# anti-HSD17B13 antibody (AA 171-270) (Biotin)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	HSD17B13	
Binding Specificity:	AA 171-270	
Reactivity:	Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HSD17B13 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human HSD17B13
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	HSD17B13
Alternative Name:	HSD17B13 (HSD17B13 Products)

## **Target Details**

Background:	Synonyms: 17-beta-HSD 13, 17-beta-hydroxysteroid dehydrogenase 13, DHB13_HUMAN,
	HMFN0376, Hsd17b13, SCDR9, SDR16C3, Short-chain dehydrogenase/reductase 9,
	UNQ497/PRO1014.
	Background: 17 beta-HSD13 (17 beta hydroxysteroid dehydrogenase type 13), also designated
	Short-chain dehydrogenase/reductase 9 (SCDR9), belongs to the 17 beta-HSD family of
	proteins, which regulate the availability of steroids within various tissues throughout the body.
	17 beta-HSD13 is a 300 amino acid secreted protein that is highly expressed in liver and is also
	detected in ovary, bone marrow, kidney, brain, lung, skeletal muscle, bladder and testis. The
	gene encoding 17 beta-HSD13 maps to chromosome 4, which houses nearly 6 % of the human
	genome and has the largest gene deserts (regions of the genome with no protein encoding
	genes) of all of the human chromosomes. Defects in some of the genes located on
	chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome,
	methylmalonic acidemia and polycystic kidney disease.
Gene ID:	345275
UniProt:	Q7Z5P4
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling
----------

Expiry Date:

12 months