antibodies -online.com





anti-NAA15 antibody (AA 121-220) (Biotin)



Go to Product page

| () | 11/0 | K\ / | iew |
|-----|--------|--------|-------------|
| | \cup | 'I V/I | $I \cap VV$ |
| | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | NAA15 | |
| Binding Specificity: | AA 121-220 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This NAA15 antibody is conjugated to Biotin | |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human NARG1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Zebrafish |
| Purification: | Purified by Protein A. |

Target Details

| Target: | NAA15 |
|-------------------|------------------------|
| Alternative Name: | NARG1 (NAA15 Products) |

Target Details

| - | | |
|---------------------|--|--|
| Background: | Synonyms: 5730450D16Rik, 6330400I15, ASTBDN, GA19, Gastric cancer antigen Ga19, | |
| | MGC29428, mNAT1, N alpha acetyltransferase 15, NatA auxiliary subunit, N terminal | |
| | acetyltransferase, N(alpha) acetyltransferase 15, NatA auxiliary subunit, N-alpha- | |
| | acetyltransferase 15, N-terminal acetyltransferase, Naa15, NAA15_HUMAN, NatA auxiliary | |
| | subunit, NATH, NMDA receptor regulated 1, NMDA receptor regulated protein 1, NMDA | |
| | receptor-regulated protein 1, Protein tubedown 1, Protein tubedown-1, Tbdn 1, TBDN100, | |
| | Transcriptional coactivator tubedown 100, tubedown-1. | |
| | Background: This gene encodes a protein of unknown function. However, similarity to proteins | |
| | in yeast and other species suggests that this protein may be an N-acetyltransferase. [provided | |
| | by RefSeq, Jul 2008] | |
| Gene ID: | 80155 | |
| UniProt: | Q9BXJ9 | |
| Application Details | | |
| Application Notes: | WB 1:300-5000 | |
| | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C for 12 months. | |
| Expiry Date: | 12 months | |
| | | |