antibodies -online.com





anti-Nesprin3 antibody (AA 201-300)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	Nesprin3 (C14orf49)
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Nesprin3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Nesprin 3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Nesprin3 (C14orf49)
Alternative Name:	Nesprin 3 (C14orf49 Products)

Target Details

Background:	Synonyms: C14orf49, Chromosome 14 open reading frame 49, FLJ25605, MGC75267, Nesp3,	
	Nesprin3, Nesprin-3, NET53, Nuclear Envelope Spectrin Repeat Protein 3, Spectrin Repeat	
	Containing, Nuclear Envelope Family Member 3, SYNE3, SYNE3_HUMAN.	
	Background: As a component of the LINC (Linker of Nucleoskeleton and Cytoskeleton) complex	
	involved in the connection between the nuclear lamina and the cytoskeleton. The	
	nucleocytoplasmic interactions established by the LINC complex play an important role in the	
	transmission of mechanical forces across the nuclear envelope and in nuclear movement and	
	positioning. Probable anchoring protein which tethers the nucleus to the cytoskeleton by	
	binding PLEC which can associate with the intermediate filament system. Plays a role in the	
	regulation of aortic epithelial cell morphology, and is required for flow-induced centrosome	
	polarization and directional migration in aortic endothelial cells	
Gene ID:	161176	
UniProt:	Q6ZMZ3	
Pathways:	Maintenance of Protein Location	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

Handling

Storage:	4 °C,-20 °C
Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months