



[Go to Product page](#)

Datasheet for ABIN6944508  
**anti-PEX5 antibody (AA 51-150) (AbBy Fluor® 555)**

### Overview

Quantity:	100 µL
Target:	PEX5
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PEX5 antibody is conjugated to AbBy Fluor® 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PEX5
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### Target Details

Target:	PEX5
Alternative Name:	PEX5 ( <a href="#">PEX5 Products</a> )

## Target Details

---

**Background:** Synonyms: FLJ50634, FLJ50721, FLJ51948, Peroxin 5, Peroxin-5, Peroxisomal biogenesis factor 5, Peroxisomal C terminal targeting signal import receptor, Peroxisomal C-terminal targeting signal import receptor, Peroxisomal targeting signal 1 receptor, Peroxisome receptor 1, pex5, PEX5\_HUMAN, PTS1 BP, PTS1 receptor, PTS1-BP, PTS1R, PXR1.

Background: The product of this gene binds to the C-terminal PTS1-type tripeptide peroxisomal targeting signal (SKL-type) and plays an essential role in peroxisomal protein import. Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. The peroxisomal biogenesis disorders are a heterogeneous group with at least 14 complementation groups and with more than 1 phenotype being observed in cases falling into particular complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of neonatal adrenoleukodystrophy (NALD), a cause of Zellweger syndrome (ZWS) as well as may be a cause of infantile Refsum disease (IRD). Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2008]

---

**Gene ID:** 5830

---

**UniProt:** [P50542](#)

---

**Pathways:** [Monocarboxylic Acid Catabolic Process](#)

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 µg/µL

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

## Handling

---

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months