

Datasheet for ABIN6944764

anti-RCAS1 antibody (AA 151-213) (AbBy Fluor® 594)



Go to Product page

| (|) | V | | rV | ĺ | 9 | V | V |
|---|---------------|----|----------|----|---|---------------|---|---|
| ' | \mathcal{I} | ٧V | <u> </u> | v | 1 | $\overline{}$ | ٧ | ٧ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | RCAS1 (EBAG9) | |
| Binding Specificity: | AA 151-213 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This RCAS1 antibody is conjugated to AbBy Fluor® 594 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RCAS1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Cow,Pig,Chicken,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | RCAS1 (EBAG9) |
|-------------------|------------------------------|
| Alternative Name: | RCAS1/EBAG9 (EBAG9 Products) |

Target Details

| Background: | Synonyms: BAG9, Cancer associated surface antigen, Cancer associated surface antigen |
|---------------------|--|
| | RCAS1, EB9, EBAG 9, Estrogen receptor binding fragment associated gene 9 protein, Estrogen |
| | receptor binding site associated antigen 9, PDAF, RCAS 1, RCAS1, Receptor binding cancer |
| | antigen expressed on SiSo cells. |
| | Background: This gene was identified as an estrogen-responsive gene. Regulation of |
| | transcription by estrogen is mediated by estrogen receptor, which binds to the estrogen- |
| | responsive element found in the 5'-flanking region of this gene. The encoded protein is a tumor- |
| | associated antigen that is expressed at high frequency in a variety of cancers. This gene may |
| | also participate in the suppression of cell proliferation and inducing apoptotic cell death |
| | through activation of interleukin-1-beta converting enzyme (ICE)-like proteases. Alternate |
| | splicing results in multiple transcript variants. A pseudogene of this gene has been defined on |
| | chromosome 10. |
| Gene ID: | 9166 |
| UniProt: | 000559 |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |

| \vdash | land | lına |
|----------|-------|------|
| | iaria | шц |

Expiry Date:

12 months