

## Datasheet for ABIN6945105

## anti-CPT1A antibody (AA 351-450) (Biotin)



Go to Product page

$\sim$				
$O_1$	<b>/</b> el	rVI	161	Λ

Quantity:	100 μL	
Target:	CPT1A	
Binding Specificity:	AA 351-450	
Reactivity:	Mouse, Rat, Pig	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CPT1A antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human CPT1A	
	KLH conjugated synthetic peptide derived from human CPT1A	
Immunogen:		
Immunogen: Isotype:	IgG	
Immunogen: Isotype: Cross-Reactivity:	IgG Mouse, Pig, Rat	
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG  Mouse, Pig, Rat  Human,Dog,Cow,Chicken	
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG  Mouse, Pig, Rat  Human,Dog,Cow,Chicken	
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification: Target Details	IgG  Mouse, Pig, Rat  Human,Dog,Cow,Chicken  Purified by Protein A.	

palmitoyltransferase 1, liver isoform, CPT I, Carnitine palmitoyltransferase 1A, carnitine 0-
palmitoyltransferase 1, liver isoform isoform 1, CPT1A_HUMAN.

Background: The mitochondrial oxidation of long-chain fatty acids is nitiated by the sequential action of carnitine palmitoyltransferase I (which is located in the outer membrane and is detergent-labile) and carnitine palmitoyltransferase II (which is located in the inner membrane and is detergent-stable), together with a carnitine-acylcarnitine translocase. CPT I is the key enzyme in the carnitine-dependent transport across the mitochondrial inner membrane and its deficiency results in a decreased rate of fatty acid beta-oxidation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].

UniProt: P50416

Pathways: AMPK Signaling, Regulation of Lipid Metabolism by PPARalpha, Feeding Behaviour,

Monocarboxylic Acid Catabolic Process

## **Application Details**

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months