antibodies -online.com





anti-USP14 antibody



Overview

Quantity:	100 μL
Target:	USP14
Reactivity:	Human, Mouse, Rat, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This USP14 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Alternative Name:

Product Details	
Immunogen:	Recombinant protein within human USP14 aa 300 to the C-terminus
Clone:	4C7
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat, Zebrafish (Danio rerio)
Purification:	Purified by Protein A.
Target Details	
Target:	USP14

Usp14 (USP14 Products)

Target Details

Storage Comment:

rarget Details	
Background:	Synonyms: Ubiquitin carboxyl-terminal hydrolase 14, Deubiquitinating enzyme 14, Ubiquitin
	thioesterase 14, Ubiquitin-specific-processing protease 14, USP14, TGT.
	Background: Proteasome-associated deubiquitinase which releases ubiquitin from the
	proteasome targeted ubiquitinated proteins. Ensures the regeneration of ubiquitin at the
	proteasome. Is a reversibly associated subunit of the proteasome and a large fraction of
	proteasome-free protein exists within the cell. Required for the degradation of the chemokine
	receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis. Serves also as a
	physiological inhibitor of endoplasmic reticulum-associated degradation (ERAD) under the nor
	stressed condition by inhibiting the degradation of unfolded endoplasmic reticulum proteins vi
	interaction with ERN1. Indispensable for synaptic development and function at neuromuscular
	junctions (NMJs).
Gene ID:	9097
UniProt:	P54578
Application Details	
Application Notes:	WB 1:300-5000
	FCM 1:20-100
	IHC-P 1:200-400
	IF(IHC-P) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 1xTBS (pH 7.4), 1 % BSA, 40 %Glycerol and 0.05 %
	Sodium Azide.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C

Store at 4°C for up to 2 weeks. For long term storage, store at -20°C in small aliquots to prevent

Handling

	freeze-thaw cycles.
Expiry Date:	12 months