

Datasheet for ABIN6945388

anti-WEE1 antibody (AA 151-250) (Cy5.5)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	WEE1
Binding Specificity:	AA 151-250
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WEE1 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human WEE1
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse
Purification:	Purified by Protein A.

Target Details

Target:	WEE1
Alternative Name:	WEE1 (WEE1 Products)

Target Details

Background:	Synonyms: DKFZp686l18166, EC 2.7.10.2, FLJ16446, Wee 1, WEE-1, WEE1 protein, Wee1 homolog, WEE1 homolog S. pombe, Wee1 like protein kinase, Wee1 tyrosine kinase, Wee1+ homolog, Wee1+ S. pombe homolog, WEE1A, Wee1A kinase, WEE1hu, WEE1 tyrosine kinase isoform 1, MGC105683, WEE1_HUMAN. Background: This gene encodes a nuclear protein, which is a tyrosine kinase belonging to the Ser/Thr family of protein kinases. Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation.
Gene ID:	7465
UniProt:	P30291
Pathways:	Cell Division Cycle, Mitotic G1-G1/S Phases, M Phase

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------