antibodies - online.com





anti-ZNF197 antibody (AA 231-330) (Biotin)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	ZNF197	
Binding Specificity:	AA 231-330	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF197 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ZNF197
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Rat,Cow,Horse,Cat
Purification:	Purified by Protein A.

Target Details

Target:	ZNF197
Alternative Name:	ZNF197 (ZNF197 Products)

Target Details

papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000			
20, Zinc finger protein with KRAB and SCAN domains 9, ZKSCAN9, ZNF166, ZNF20. Background: This gene product belongs to the zinc finger protein superfamily, members of which are regulatory proteins characterized by nucleic acid-binding zinc finger domains. The encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR AUUUA motifs, which are often a hailmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt. 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use. This product contains ProClin: a POISONOUS AND HAZAROOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 morths.	Background:	Synonyms: D3S1363E, OTTHUMP00000164119, P18, VHL associated KRAB A domain	
Background: This gene product belongs to the zinc finger protein superfamily, members of which are regulatory proteins characterized by nucleic acid-binding zinc finger domains. The encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq. Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Clycerol Preservative: ProClin Precaution of Use. This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		containing protein, VHLaK, Zinc finger protein 166, Zinc finger protein 197, Zinc finger protein	
which are regulatory proteins characterized by nucleic acid-binding zinc finger domains. The encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.		20, Zinc finger protein with KRAB and SCAN domains 9, ZKSCAN9, ZNF166, ZNF20.	
encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000		Background: This gene product belongs to the zinc finger protein superfamily, members of	
box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq. Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-P 1:00-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should b handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		which are regulatory proteins characterized by nucleic acid-binding zinc finger domains. The	
AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyro papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated	
papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		box (KRAB) domain, and a SCAN box. This transcript turns over rapidly and contains 3' UTR	
alternatively spliced transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008] Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should by handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		AUUUA motifs, which are often a hallmark of rapid turnover. It is overexpressed in some thyroid	
Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		papillary carcinomas. This gene is located in a cluster of zinc finger genes at 3p21. Two	
Gene ID: 10168 UniProt: 014709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		alternatively spliced transcripts encoding different isoforms have been described. [provided by	
UniProt: O14709 Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		RefSeq, Jul 2008]	
Application Details Application Notes: WB 1:300-5000 IHC-F 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.	Gene ID:	10168	
Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.	UniProt:	014709	
IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.	Application Details		
Handling For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.	Application Notes:	WB 1:300-5000	
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C for 12 months.		IHC-P 1:200-400	
Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 at 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.		IHC-F 1:100-500	
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Restrictions:	For Research Use only	
Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ard 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Handling		
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Format:	Liquid	
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Concentration:	1 μg/μL	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: Storage Comment: Store at -20°C for 12 months.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.		50 % Glycerol.	
handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Preservative:	ProClin	
Storage: -20 °C Storage Comment: Store at -20 °C for 12 months.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
Storage Comment: Store at -20°C for 12 months.		handled by trained staff only.	
	Storage:	-20 °C	
Expiry Date: 12 months	Storage Comment:	Store at -20°C for 12 months.	
	Expiry Date:	12 months	