

[Go to Product page](#)

Datasheet for ABIN6951197

TRKB ELISA Kit

Overview

Quantity:	96 tests
Target:	TRKB (NTRK2)
Binding Specificity:	pTyr816
Reactivity:	Human, Rat, Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human, Mouse and Rat Phospho-TrkB (Tyr816) ELISA Kit. This assay semi-quantitatively measures TrkB phosphorylated at Tyrosine-816 in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibodies provided in this kit recognizes human, mouse, and rat TrkB phosphorylated at Tyrosine-816.
Characteristics:	<ul style="list-style-type: none">• Rapidly measure phosphorylated protein in lysates• Screen numerous different cell lysates without performing a Western Blot analysis• Minimal hands-on time, convenient, and non-radioactive material
Components:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer• Anti-Phospho Antibody

Product Details

- HRP-Conjugated Secondary Antibody
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 μ L to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: TRKB (NTRK2)

Alternative Name: TrkB ([NTRK2 Products](#))

Gene ID: 4915

Pathways: [RTK Signaling](#), [Neurotrophin Signaling Pathway](#), [cAMP Metabolic Process](#), [Skeletal Muscle Fiber Development](#), [Feeding Behaviour](#), [Dicarboxylic Acid Transport](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 μ L of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 $^{\circ}$ C.
4. Add 100 μ L of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 μ L of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Application Details

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months