

## Datasheet for ABIN6951489

# **E-cadherin ELISA Kit**



### Overview

Quantity:	96 tests
Target:	E-cadherin (CDH1)
Reactivity:	Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Application.	LLIOA
Product Details	
Purpose:	Rat E-Cadherin ELISA Kit.
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Characteristics:	<ul> <li>Strip plates and additional reagents allow for use in multiple experiments</li> <li>Quantitative protein detection</li> <li>Establishes normal range</li> <li>The best products for confirmation of antibody array data</li> </ul>
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Stop Solution</li> <li>Assay Diluent(s)</li> <li>Lyophilized Standard</li> <li>Biotinylated Detection Antibody</li> <li>Streptavidin-Conjugated HRP</li> <li>TMB One-Step Substrate</li> </ul>

### **Product Details**

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- Log-log graph paper or computer and software for ELISA data analysis

## Target Details

Target:	E-cadherin (CDH1)
Alternative Name:	E-Cadherin (CDH1 Products)
Gene ID:	83502
UniProt:	Q9R0T4
Pathways:	WNT Signaling, Sensory Perception of Sound, Cell-Cell Junction Organization, Tube Formation

## **Application Details**

Optimal working dilution should be determined by the investigator.  Pre-coated  1. Prepare all reagents, samples and standards as instructed in the manual.
1. Prepare all reagents, samples and standards as instructed in the manual
r. i repare an reagents, samples and standards as instructed in the mandal.
2. Add 100 μl of standard or sample to each well.
3. Incubate 2.5 h at RT or O/N at 4°C.
4. Add 100 μl of prepared biotin antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µl of prepared Streptavidin solution to each well.
7. Incubate 45 min at RT.
8. Add 100 µl of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μl of Stop Solution to each well.
11. Read at 450 nm immediately.
For Research Use only
6 months