# antibodies -online.com







# **HMOX1 ELISA Kit**



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	96 tests
Target:	HMOX1
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

1 1		
Draduat Dataila		
Product Details		
Purpose:	Human HSP32/HO-1/HMOX1 ELISA Kit.	
Sample Type:	Cell Culture Supernatant, Cell Samples, Plasma, Serum, Tissue Lysate	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Characteristics:	Strip plates and additional reagents allow for use in multiple experiments	
	Quantitative protein detection	
	Establishes normal range	
	The best products for confirmation of antibody array data	
Components:	Pre-Coated 96-well Strip Microplate	
	Wash Buffer	
	Stop Solution	
	Assay Diluent(s)	
	Lyophilized Standard	
	Biotinylated Detection Antibody	
	Streptavidin-Conjugated HRP	
	TMB One-Step Substrate	

### **Product Details**

#### Material not included:

- · Distilled or deionized water
- Precision pipettes to deliver 2 μl to 1 μl volumes
- Adjustable 1-25 µl pipettes for reagent preparation
- 100 µl and 1 liter graduated cylinders
- Tubes to prepare standard and sample dilutions
- · Absorbent paper
- Microplate reader capable of measuring absorbance at 450nm
- · Log-log graph paper or computer and software for ELISA data analysis

## Target Details

Target:	HMOX1	
Alternative Name:	HSP32 (HO-1/HMOX1) (HMOX1 Products)	
Gene ID:	3162	
UniProt:	P09601	
Pathways:	Transition Metal Ion Homeostasis, Regulation of Leukocyte Mediated Immunity, Positive	
	Regulation of Immune Effector Process, Production of Molecular Mediator of Immune	
	Response, SARS-CoV-2 Protein Interactome	

## **Application Details**

Application Notes:	Optimal working dilution should be determined by the investigator.	
Plate:	Pre-coated	
Protocol:	1. Prepare all reagents, samples and standards as instructed in the manual.	
	2. Add 100 µl of standard or sample to each well.	
	3. Incubate 2.5 h at RT or O/N at 4°C.	
	4. Add 100 μl of prepared biotin antibody to each well.	
	5. Incubate 1 h at RT.	
	6. Add 100 µl of prepared Streptavidin solution to each well.	
	7. Incubate 45 min at RT.	
	8. Add 100 µl of TMB One-Step Substrate Reagent to each well.	
	9. Incubate 30 min at RT.	
	10. Add 50 μl of Stop Solution to each well.	
	11. Read at 450 nm immediately.	
Restrictions:	For Research Use only	

1 1		
$\blacksquare$	$\alpha \alpha \alpha$	lina
	land	1111(
	I G I G	11119

Expiry Date:

6 months