
antibodies - online.com







anti-tau antibody (Biotin)

Images



Overview

| Quantity: | 100 μg |
|--------------|---|
| Target: | tau |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This tau antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Dot Blot (DB) |

Product Details


| Immunogen: | Human Recombinant Tau441 (2N4R), P301S mutant Protein Pre-formed Fibrils |
|-------------------|--|
| Clone: | 1D5 |
| Isotype: | lgG1 |
| Specificity: | Expressed in neurons.,Detects Multiple Bands. Antibody detects monomer under denaturing conditions but preferentially detects fibril under native conditions (dot blot). |
| Cross-Reactivity: | Human, Mouse |
| Purification: | Protein G Purified |


Target Details


Target: tau

Target Details

| Alternative Name: | Tau (tau Products) |
|---------------------|--|
| Background: | Alzheimer's Disease (AD) is the most common neurodegenerative disease, affecting 10 % of seniors over the age of 65 (1). It was named after Alois Alzheimer, a German scientist who discovered tangled bundles of fibrils where neurons had once been in the brain of a deceased patient in 1907 (2). Tau (tubulin-associated unit) is normally located in the axons of neurons where it stabilizes microtubules. Tauopathies such as AD are characterized by neurofibrillary tangles containing hyperphosphorylated tau fibrils (3). There are six isoforms of tau in the adul human brain: three with four repeat units (4R) and three with three repeat units (3R) (4). 2N4R, or Tau-441 is the full length tau protein. P301S is a mutation encoded by exon 10 (4) that impairs the ability of tau to assemble microtubules (5). |
| NCBI Accession: | NP_005901 |
| UniProt: | P10636 |
| Application Details | |
| Application Notes: | WB (1:1000)optimal dilutions for assays should be determined by the user. |
| Comment: | A 1:1000 dilution of ABIN6952089 was sufficient for detection of Tau 2N4R P301S Fibril in 20 ug of Mouse Brain by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.09 % Sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Conjugated antibodies should be stored at 4°C |







Immunofluorescence

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Tau Monoclonal Antibody, Clone 1D5 (ABIN6952089). Tissue: iPSC-derived neurons. Species: Human. Fixation: 4 % PFA. Primary Antibody: Mouse Anti-Tau Monoclonal Antibody (ABIN6952089) at 1:100 for Overnight at 4 °C. Counterstain: DAPI at 1:5000 for 5 minutes at RT in the dark. Magnification: 40X. Courtesy of: Francesco Paonessa.

Western Blotting

Image 2. Western Blot analysis of Human Breast Cancer Cell line and Mouse Brain showing detection of Tau protein using Mouse Anti-Tau Monoclonal Antibody, Clone 1D5 (ABIN6952089). Lane 1: MW Marker. Lane 2: Human T-47d (10 μg). Lane 3: Mouse Brain (20 μg). Block: 5 % Skim Milk powder in TBST. Primary Antibody: Mouse Anti-Tau Monoclonal Antibody (ABIN6952089) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:5000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT.

Dot Blot

Image 3. Dot Blot analysis using Mouse Anti-Tau Monoclonal Antibody, Clone 1D5 (ABIN6952089). Tissue: Recombinant Protein. Species: Human. Primary Antibody: Mouse Anti-Tau Monoclonal Antibody (ABIN6952089) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:5000 for 1 hour at RT with shaking.