

Datasheet for ABIN6952258

**Fc epsilon RI/FCER1A Protein (AA 26-204) (Fc Tag)**[Go to Product page](#)**1** Image

## Overview

Quantity:	100 µg
Target:	Fc epsilon RI/FCER1A (FCER1A)
Protein Characteristics:	AA 26-204
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Fc epsilon RI/FCER1A protein is labelled with Fc Tag.

## Product Details

Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

## Target Details

Target:	Fc epsilon RI/FCER1A (FCER1A)
Alternative Name:	Fc epsilon RI alpha ( <a href="#">FCER1A Products</a> )
Background:	High affinity immunoglobulin epsilon receptor subunit alpha (FCER1A) is also known as Fc-epsilon RI-alpha (FcERI), IgE Fc receptor subunit alpha, FCE1A. FCER1A contains two Ig-like (immunoglobulin-like) domains. FCER1A binds to the Fc region of immunoglobulins epsilon and is a high affinity receptor. FCER1A is responsible for initiating the allergic response, which binding of allergen to receptor-bound IgE leads to cell activation and the release of mediators (such as histamine) responsible for the manifestations of allergy. The same receptor also

Target Details

	induces the secretion of important lymphokines. FCER1A plays a central role in allergic disease, coupling allergen and mast cell to initiate the inflammatory and immediate hypersensitivity responses that are characteristic of disorders such as hay fever and asthma.
Molecular Weight:	47.3 kDa
NCBI Accession:	<a href="#">NP_001992</a>
Pathways:	<a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Positive Regulation of Immune Effector Process</a>


Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Buffer:	Tris with Glycine, Arginine and NaCl, pH 7.5
Storage:	-20 °C

Images



**SDS-PAGE**

**Image 1.** Human Fc epsilon RI alpha, Fc Tag (BLI verified) on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .