

Datasheet for ABIN6952338

anti-Histone 3 antibody (H3K27me3)



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	50 μg
Target:	Histone 3 (H3)
Binding Specificity:	H3K27me3
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunofluorescence (IF), ELISA, Chromatin Immunoprecipitation (ChIP), Dot Blot (DB), ChIP DNA-Sequencing (ChIP-seq), Cleavage Under Targets and Tagmentation (CUT&Tag)

Product Details

Immunogen:	synthetic peptide
Specificity:	Polyclonal antibody raised in rabbit against against histone H3, trimethylated at lysine 27 (H3K27me3), using a KLH-conjugated synthetic peptide.
Cross-Reactivity:	Arabidopsis, Cow (Bovine), Fruit Fly (Drosophila melanogaster), Mouse (Murine), Pig (Porcine), Rat (Rattus), Schistosoma japonicum, Zebrafish (Danio rerio)
Purification:	Peptide affinity purified
Grade:	ChIP-seq Grade

Target Details

Target: Histone 3 (H3)

Target Details

Alternative Name:	Histone 3 (H3 Products)
Background:	Histones are the main constituents of the protein part of chromosomes of eukaryotic cells.
	They are rich in the amino acids arginine and lysine and have been greatly conserved during
	evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each
	class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one
	octameric nucleosome. Histone tails undergo numerous post-translational modifications,
	which either directly or indirectly alter chromatin structure to facilitate transcriptional activation
	or repression or other nuclear processes. In addition to the genetic code, combinations of the
	different histone modifications reveal the so-called "histone code". Histone methylation and
	demethylation is dynamically regulated by respectively histone methyl transferases and histone
	demethylases. Trimethylation of histone H3K27 is associated with gene repression.
UniProt:	P68431

Application Details

Precaution of Use:

Storage:

Application Notes:	ChIP/ChIP-seq 1 μg:ChIP
	ELISA 1:5,000
	Dot Blot 1:20,000
	WB 1:1,000
	IF 1:500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1.6 μg/μL
Buffer:	PBS, 0.05 % azide, 0.05 % ProClin300
	1 20, 0.00 % delac, 0.00 % 1 100 mileo

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

should be handled by trained staff only.

-20 °C