

## Datasheet for ABIN6961764 anti-KLRG1 antibody



Overview

Quantity:	100 µg
Target:	KLRG1
Reactivity:	Mouse
Host:	Golden Syrian Hamster
Clonality:	Monoclonal
Conjugate:	This KLRG1 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunoprecipitation (IP)

## Product Details

Clone:	2F1
Isotype:	IgG
Purification:	This monoclonal antibody preparation was purified from tissue culture supernatant via affinity chromatography. For In Vivo Ready <sup>™</sup> (IVR) products, each preparation is also evaluated for endotoxin levels using the LAL assay. It is recommended to store the product undiluted at 4°C. Do not freeze.

## Target Details

Target:	KLRG1
Alternative Name:	KLRG1 (KLRG1 Products)
Background:	The 2F1 antibody reacts with mouse KLRG1 (Killer cell Lectin-like Receptor G1). This 30-38 kDa
	homodimeric receptor may be expressed by activated, mature NK cells and by

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6961764 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

	effector/memory T cells, with potentially different roles in each cell type. KLRG1 can regulate, in
	an inhibitory fashion, the development and effector functions of NK cells, and is often cited as a
	senescence or terminal differentiation marker for T cells. Ligands for KLRG1 include members
	of the cadherin family of adhesion molecules, specifically N-Cadherin, E-Cadherin, and R-
	Cadherin. These interactions may induce bidirectional, immunosuppressive signaling in both
	KLRG- and Cadherin-expressing cells. A more recently identified role for KLRG1-Cadherin
	signaling in tissue organization, e.g. in cardiac angiogenesis, expands the function of these
	interactions beyond immunosuppression of immune cells. (Bouchentouf et al. 2010. J.
	Immunol. 185: 7014-7025).The 2F1 antibody may be used as a phenotypic marker for KLRG1 in
	mouse, frequently in combination with Anti-Mouse CD127 antibody (clone A7R34), for
	identification of effector T cell populations.
Gene ID:	50928
UniProt:	088713
Application Details	
Application Notes:	This purified format is guaranteed to be >90 % pure as determined by SDS-PAGE analysis.
Comment:	0.5 mg/mL
Restrictions:	For Research Use only
Handling	

Buffer:	10 mM NaH2PO4, 150 mM NaCl, 0.09 % Sodium azide, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	2-8°C
Expiry Date:	12 months