

Datasheet for ABIN6963781

anti-Interferon gamma antibody (PromoFluor 647 Premium)



Image



(11 /	\sim	× 1	i.	\sim		
\cup	٧	Н	rv	I	H	٧	1

Quantity:	100 tests	
Target:	Interferon gamma (IFNG)	
Reactivity:	Hispid Cotton Rat	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This Interferon gamma antibody is conjugated to PromoFluor 647 Premium	
Application:	Flow Cytometry (FACS)	

Product Details

Immunogen:	Recombinant cotton rat IFN-gamma	
Clone:	MT417	
Isotype:	lgG1	
Specificity:	Native and recombinant IFN-γ	
Purification:	Purified from in vitro cultures by protein G affinity chromatography.	

Target Details

ulation of Immune Effector Process, Production of

Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy

Application Details

Application Notes:


Detection of cotton rat IFN- γ producing cells by flow cytometry. 5 μ L is recommended for staining of 1 million cells in a total volume of 50 μ L. Detection of IFN- γ by flow cytometry in viable cotton rat spleen cells. Cells were stimulated for 16 hours in the presence of PMA/ ionomycin and Brefeldin A. Cells were then fixed and permeabilized using 4 % paraformaldehyde and saponin, and subsequently stained with MT417-PF647P (solid line). Matched isotype control antibody (dashed line). The histogram derives from gated events of typical lymphocyte characteristics in forward and side light scatter. Flow cytometry was performed on a BD FACSVerse system.

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Buffer:	in PBS with 0.2 % BSA and 0.09 % sodium azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store product at 4-8°C or frozen at -20°C or below. Avoid repeated freezing/thawing.	
Expiry Date:	18 months	



Flow Cytometry

Image 1. Detection of IFN-γ by flow cytometry in viable cotton rat spleen cells. Stimulated, fixed and permeabilized cells were stained with anti-cotton rat mAb MT417-PF647P (solid line). Matched isotype control antibody (dashed line). Flow cytometry was performed on a BD FACSVerse system.