

Datasheet for ABIN6963790

anti-Interferon gamma antibody (FITC)





Go to Product page

| _ | | | | | | |
|---|-----|----------|----|----|----|----|
| | 1// | Д | rv | 16 | ٦/ | ٨ |
| U | W | \vdash | ΙV | Ιt | ٦, | /V |

| Quantity: | 100 tests |
|--------------|------------------------------------------------------|
| Target: | Interferon gamma (IFNG) |
| Reactivity: | Pig |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This Interferon gamma antibody is conjugated to FITC |
| Application: | Flow Cytometry (FACS) |

Product Details

| Immunogen: | Recombinant porcine IFN-gamma |
|---------------|-----------------------------------------------------------------------|
| Clone: | P2C11 |
| Isotype: | lgG2a |
| Specificity: | Native and recombinant porcine IFN-γ |
| Purification: | Purified from in vitro cultures by protein G affinity chromatography. |

Target Details

| Target: | Interferon gamma (IFNG) |
|-------------------|--------------------------------------------------------------------------------------------|
| Alternative Name: | IFNG (IFNG Products) |
| Gene ID: | 396991 |
| Pathways: | Interferon-gamma Pathway, Cellular Response to Molecule of Bacterial Origin, Regulation of |

Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, ER-Nucleus Signaling, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy

Application Details

Application Notes:


Detection of IFN- γ producing cells by flow cytometry. 5 µL is recommended for staining of 1 million cells in a total volume of 50 µL. Detection of IFN- γ by flow cytometry in viable porcine PBMC. Cells were stimulated for 16 hours in the presence of PMA/ionomycin and Brefeldin A. Cells were then fixed and permeabilized using 4 % paraformaldehyde and saponin, and subsequently stained with CD3- PE and P2C11-FITC (right panel). The left panel depicts cells stained with a matched isotype control antibody. The dot plots derive from gated events of typical lymphocyte characteristics in forward and side light scatter. Flow cytometry was performed on a BD FACSVerse system.

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|------------------------------------------------------------------------------------------------------------------------|
| Buffer: | in PBS with 0.2 % BSA and 0.09 % sodium azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store product at 4-8°C or frozen at -20°C or below. Avoid repeated freezing/thawing. |
| Expiry Date: | 18 months |



Flow Cytometry

Image 1. Detection of IFN- γ by flow cytometry in viable porcine PBMC. Stimulated, fixed and permeabilized cells were stained with anti-porcine CD3-PE and anti-porcine IFN- γ mAb P2C11-FITC (right panel). Matched isotype control antibody (left panel). Flow cytometry was performed on a BD FACSVerse system.