

Datasheet for ABIN6964383 CD1d Protein (Fc Tag)



Overview	
Overview	
Quantity:	100 µg
Target:	CD1d (CD1D)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD1d protein is labelled with Fc Tag.
Product Details	
Sequence:	CD1D (Gln20-Ser301)+hFc(GLU99-ALA330)
Characteristics:	Tagged Protein: C-Human Fc tag
Purification:	affinity purification
Target Details	
Target:	CD1d (CD1D)
Alternative Name:	CD1D (CD1D Products)
Background:	Synonymes: CD1A, R3, R3G1
	Description: This gene encodes a divergent member of the CD1 family of transmembrane
	glycoproteins, which are structurally related to the major histocompatibility complex (MHC)
	proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the
	presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The
	human genome contains five CD1 family genes organized in a cluster on chromosome 1. The

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6964383 | 05/14/2025 | Copyright antibodies-online. All rights reserved.

	CD1 family members are thought to differ in their cellular localization and specificity for
	particular lipid ligands. The protein encoded by this gene localizes to late endosomes and
	lysosomes via a tyrosine-based motif in the cytoplasmic tail. Two transcript variants encoding
	different isoforms have been found for this gene. [provided by RefSeq, Jan 2016]
Molecular Weight:	56.54 kDa
UniProt:	P15813
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Handling Format:	Lyophilized
	Lyophilized Reconstitute with deionized water
Format:	
Format: Reconstitution:	Reconstitute with deionized water
Format: Reconstitution:	Reconstitute with deionized water Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants
Format: Reconstitution: Buffer:	Reconstitute with deionized water Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Format: Reconstitution: Buffer: Preservative:	Reconstitute with deionized water Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization. Without preservative