



[Go to Product page](#)

Datasheet for ABIN6972985

## CD36 Protein (CD36) (AA 30-439) (His tag,AVI tag,Biotin)

### 2 Images

#### Overview

Quantity:	200 µg
Target:	CD36
Protein Characteristics:	AA 30-439
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD36 protein is labelled with His tag,AVI tag,Biotin.

#### Product Details

Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	Biotinylated Human CD36 / SR-B3 Protein, His,Avitag™
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

#### Target Details

Target:	CD36
Alternative Name:	CD36 ( <a href="#">CD36 Products</a> )
Background:	CD36 (Cluster of Differentiation 36) is also known as platelet membrane glycoprotein IV (GPIV),

## Target Details

---

fatty acid translocase (FAT), thrombospondin receptor, collagen receptor, and scavenger receptor class B, member 3 (SRB3), is a member of the class B scavenger receptor family of cell surface proteins. The human CD36 gene encodes a single chain 472 amino acid residue protein containing both an N- and a C-terminal cytoplasmic tail and an extracellular loop. CD36 is found on platelets, erythrocytes, monocytes, differentiated adipocytes, mammary epithelial cells, spleen cells and some skin microdermal endothelial cells. CD36 is a multiligand pattern recognition receptor that interacts with a large number of structurally dissimilar ligands, including long chain fatty acid (LCFA), advanced glycation end products (AGE), thrombospondin-1, oxidized low-density lipoproteins (oxLDLs), high density lipoprotein (HDL), phosphatidylserine, apoptotic cells, beta-amyloid fibrils (fA $\beta$ ), collagens I and IV, and Plasmodium falciparum infected erythrocytes. CD36 is required for the anti-angiogenic effects of thrombospondin1 In the corneal neovascularization assay. On binding a ligand the protein and ligand are internalized. This internalization is independent of macropinocytosis and occurs by an actin dependent mechanism requiring the activation Src-family kinases, JNK and Rho-family GTPases. CD36 ligands have also been shown to promote sterile inflammation through assembly of a Toll-like receptor 4 and 6 heterodimer.

---

Molecular Weight: 50.2 kDa

---

NCBI Accession: [NP\\_001001547](#)

---

Pathways: [TLR Signaling](#), [Peptide Hormone Metabolism](#), [Response to Growth Hormone Stimulus](#), [Activation of Innate immune Response](#), [Cellular Response to Molecule of Bacterial Origin](#), [Regulation of Lipid Metabolism by PPARAlpha](#), [Positive Regulation of Immune Effector Process](#), [Production of Molecular Mediator of Immune Response](#), [Hepatitis C](#), [Toll-Like Receptors Cascades](#), [Lipid Metabolism](#), [S100 Proteins](#)

## Application Details

---

Comment: Ready-to-use Avitag™ biotinylated protein:

The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA.

This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized

## Application Details

on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.

Restrictions: For Research Use only



## Handling

Format: Lyophilized

Buffer: PBS, pH 7.4

Storage: -20 °C

## Images



### ELISA

**Image 1.** Immobilized Human R-Spondin 1, His Tag (ABIN2181684, ABIN2181683) at 5 µg/mL (100 µL/well) can bind Biotinylated Human CD36, His,Avitag (ABIN6972985) with a linear range of 0.313-1.25 µg/mL (QC tested).

### SDS-PAGE

**Image 2.** Biotinylated Human CD36, His,Avitag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 %.