

Datasheet for ABIN6973192

PD-1 Protein (AA 25-167) (His tag,Fc Tag,AVI tag,Biotin)



[Go to Product page](#)

3 Images

Overview

Quantity:	200 µg
Target:	PD-1 (PDCD1)
Protein Characteristics:	AA 25-167
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This PD-1 protein is labelled with His tag,Fc Tag,AVI tag,Biotin.

Product Details

Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	Biotinylated Human PD-1 / PDCD1 Protein, Fc,Avitag™,His Tag (MALS verified)
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	PD-1 (PDCD1)
Alternative Name:	PD-1 (PDCD1 Products)
Background:	Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I

Target Details

membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN- γ . PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN- γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN- γ secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer.

Molecular Weight: 45.5 kDa

NCBI Accession: [NP_005009](#)

Pathways: [Cancer Immune Checkpoints](#)

Application Details

Application Notes: MALS verified

Comment: Ready-to-use Avitag™ biotinylated protein:
The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA.

This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.

Restrictions: For Research Use only

Handling




Format: Lyophilized

Handling

Buffer: PBS, pH 7.4

Storage: -20 °C

Images



ELISA

Image 1. Immobilized Human PD-L2, Fc Tag (HPLC verified) (ABIN2181601,ABIN2749238) at 5 µg/mL (100 µL/well) can bind Biotinylated Human PD-1, Fc,Avitag,His Tag (ABIN6973192) with a linear range of 10-40 ng/mL (Routinely tested).

SDS-PAGE

Image 2. Biotinylated Human PD-1, Fc,Avitag,His Tag on under reducing (R) and ing (NR) conditions. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .

ELISA

Image 3. Immobilized Human PD-L1, Fc Tag (Hied) (ABIN2181596,ABIN2181595) at 5 µg/mL (100 µL/well) can bind Biotinylated Human PD-1, Fc,Avitag,His Tag (ABIN6973192) with a linear range of 10-156 ng/mL (QC tested).