

Datasheet for ABIN6973209

PVRIG Protein (AA 41-171) (Fc Tag,AVI tag,Biotin)**3** Images[Go to Product page](#)

Overview

Quantity:	200 µg
Target:	PVRIG
Protein Characteristics:	AA 41-171
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This PVRIG protein is labelled with Fc Tag,AVI tag,Biotin.

Product Details

Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	Biotinylated Human PVRIG Protein, Fc,Avitag™ (HPLC verified)
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	PVRIG
Alternative Name:	PVRIG (PVRIG Products)
Background:	Human PVRIG (poliovirus receptor related immunoglobulin domain-containing protein), also

Target Details

known as CD112 receptor (CD112R), is an approximately 34 kDa single transmembrane protein in the poliovirus receptor-like protein (PVR) family. The CD112R gene encodes a putative single transmembrane protein, which is composed of a single extracellular IgV domain, one transmembrane domain, and a long intracellular domain. Notably, the intracellular domain of phatases. The extracellular domain sequence of human and mouse CD112R have 65.3 % similarity. CD112R may act as a coinhibitory receptor that suppresses T-cell receptor-mediated signals.

Molecular Weight: 41.9 kDa

NCBI Accession: [NP_076975](#)

Application Details

Application Notes: HPLC verified

Comment: Ready-to-use Avitag™ biotinylated protein:

The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA.

This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.




Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: PBS, pH 7.4

Storage: -20 °C



ELISA

Image 1. Immobilized Human Nectin-2, Mouse IgG2a Fc Tag (ABIN6386453,ABIN6388251) at 10 µg/mL (100 µ L/well) can bind Biotinylated Human PVRIG, Fc,Avitag (HPLC verified) (ABIN6973209) with a linear range of 5-40 ng/mL (QC tested).

SDS-PAGE

Image 2. Biotinylated Human PVRIG, Fc,Avitag™ (HPLC verified) on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 % .

High Pressure Liquid Chromatography

Image 3. The purity of Biotinylated Human PVRIG, Fc,Avitag,His Tag (ABIN6973209) was greater than 90 % as determined by .