

Datasheet for ABIN6973252

SARS-CoV-2 Spike S1 Protein (NTD) (His tag,AVI tag,Biotin)[Go to Product page](#)**2** Images

Overview

Quantity:	200 µg
Target:	SARS-CoV-2 Spike S1
Protein Characteristics:	NTD
Origin:	SARS Coronavirus-2 (SARS-CoV-2)
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This SARS-CoV-2 Spike S1 protein is labelled with His tag,AVI tag,Biotin.

Product Details

Sequence:	AA 16-318
Specificity:	Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.
Characteristics:	Biotinylated S1 protein NTD, His,Avitag™ is expressed from human 293 cells (HEK293). It contains AA Val 16 - Phe 318 (Accession # QHD43416.1).
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	SARS-CoV-2 Spike S1
---------	---------------------

Target Details



Alternative Name:	SARS-CoV-2 S1 protein (SARS-CoV-2 Spike S1 Products)
Target Type:	Viral Protein
Background:	It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.
Molecular Weight:	38.0 kDa

Application Details

Comment:	Ready-to-use Avitag™ biotinylated protein: The product is exclusively produced using the Avitag™ technology. Briefly, a unique 15 amino acid peptide, the Avi tag, is introduced into the recombinant protein during expression vector construction. The single lysine residue in the Avi tag is enzymatically biotinylated by the E. Coli biotin ligase BirA. This single-point enzymatic labeling technique brings many advantages for commonly used binding assays. The biotinylation happens on the lysine residue of Avi tag, and therefore does NOT interfere with the target protein's natural binding activities. In addition, when immobilized on an avidin-coated surface, the protein orientation is uniform because the position of the Avi tag in the protein is precisely controlled.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	PBS, pH 7.4
Storage:	-20 °C



ELISA

Image 1. Immobilized A-CoV-2 Spike NTD Antibody, Chimeric mAb at 1 µg/mL (100 µL/well) can bind Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag (ABIN6973252) with a linear range of 1-25 ng/mL (QC tested).

SDS-PAGE

Image 2. Biotinylated SARS-CoV-2 S1 protein NTD, His,Avitag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 % .