# antibodies -online.com





# anti-NAT15 antibody (AA 1-100) (Alexa Fluor 350)



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	NAT15
Binding Specificity:	AA 1-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NAT15 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

#### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HAT4
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	NAT15
Alternative Name:	HAT4 (NAT15 Products)

## **Target Details**

Synonyms: HAT4, NAT15, N-alpha-acetyltransferase 60, Histone acetyltransferase type B	
protein 4, N-acetyltransferase 15, NatF catalytic subunit, NAA60, UNQ2771/PRO7155	
Background: Histone acetyltransferase localized in the Golgi apparatus that mediates	
acetylation of free histone H4, thereby facilitating nucleosome assembly. Has a preference for	
free histone H4 'Lys-20'(H4K20ac), 'Lys-79'(H4K79ac) and 'Lys-91' (H4K91ac). Also displays	
alpha (N-terminal) acetyltransferase activity towards a range of N-terminal sequences including	
those starting with Met-Lys, Met-Val, Met-Ala and Met-Met. Required for normal chromosomal	
segregation during anaphase.	
79903	
Q9H7X0	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
handled by trained staff only.  -20 °C	